

# Misspecification-Robust Inference in Linear Asset-Pricing Models with Irrelevant Risk Factors

**Nikolay Gospodinov**  
Federal Reserve Bank of Atlanta

**Raymond Kan**  
University of Toronto

**Cesare Robotti**  
Imperial College London

This paper shows that in misspecified models with risk factors that are uncorrelated with the test asset returns, the conventional inference methods tend to erroneously conclude, with high probability, that these factors are priced. Our proposed model selection procedure, which is robust to identification failure and potential model misspecification, restores the standard inference and proves to be effective in eliminating factors that do not improve the model's pricing ability. Applying our methodology to several popular asset-pricing models suggests that only the market and book-to-market factors appear to be priced, while the statistical evidence on the pricing ability of many macroeconomic factors is rather weak. (*JEL* G12, C12, C52)

The main objective of this paper is to raise the awareness of applied researchers about the pitfalls of incorrectly assuming correct specification and identification of asset-pricing models. We show that in the presence of misspecification and lack of identification, the finite-sample distributions of the statistics of interest can depart substantially from the standard asymptotic approximations developed under the assumption of correctly specified and fully

---

We thank the Editor (Andrew Karolyi), two anonymous referees, Daniel Andrei, Anton Braun, Chu Zhang, Guofu Zhou, seminar participants at the Federal Reserve Bank of Atlanta, Imperial College London, Queen's University, Singapore Management University, and University of Georgia, and participants at the 3rd Annual CIRPÉE Applied Financial Time Series Workshop at HEC (Montreal), Mathematical Finance Days 2012, the 2013 Northern Finance Association Meetings, and the 2013 SoFIE Conference for helpful discussions and comments. Gospodinov gratefully acknowledges financial support from Fonds de recherche sur la société et la culture (FQRSC), Institut de Finance Mathématique de Montréal (IFM2), and the Social Sciences and Humanities Research Council of Canada. Kan gratefully acknowledges financial support from the Social Sciences and Humanities Research Council of Canada and the National Bank Financial of Canada. The views expressed here are the authors' and not necessarily those of the Federal Reserve Bank of Atlanta or the Federal Reserve System. Supplementary data can be found on the *Review of Financial Studies* web site. Send correspondence to Cesare Robotti, Imperial College Business School, Imperial College London, Tanaka Building, South Kensington Campus, London SW7 2AZ, United Kingdom; telephone: +44 20 7594 2682. E-mail: c.robotti@imperial.ac.uk.

© The Author 2014. Published by Oxford University Press on behalf of The Society for Financial Studies.  
All rights reserved. For Permissions, please e-mail: journals.permissions@oup.com.  
doi:10.1093/rfs/hht135

Advance Access publication January 29, 2014

identified models. In general, ignoring possible model misspecification and identification failure tends to result in an overly positive assessment of the pricing performance of the asset-pricing model and the individual risk factors. To address these problems, we propose a new approach, which is robust to potential model misspecification and lack of identification, to selecting risk factors and determining whether they are priced.

It is now widely documented that misspecification is an inherent feature of many asset-pricing models, and reliable statistical inference crucially depends on its robustness to potential model misspecification. Kan and Robotti (2008, 2009), Kan, Robotti, and Shanken (2013), and Gospodinov, Kan, and Robotti (2013) show that by ignoring model misspecification, one can mistakenly conclude that a risk factor is priced when, in fact, it does not contribute to the pricing ability of the model. While these papers provide a general statistical framework for inference, evaluation and comparison of potentially misspecified asset-pricing models (see also Ludvigson 2013), the misspecification-robust inference is developed under the assumption that the covariance matrix of asset returns and risk factors is of full column rank—that is, the parameters of interest in these models are identified. Importantly, the issues with statistical inference under potential model misspecification become particularly acute when the pricing model includes factors that exhibit very low correlations with the returns on the test assets, which may jeopardize the identifiability of the model parameters.

In this paper, we further generalize the setup in the articles mentioned above to allow for possible identification failure in a stochastic discount factor (SDF) framework. We show that in the extreme case of model misspecification with one or more “useless” factors (i.e., factors that are independent of the asset returns), the identification condition fails and the validity of the statistical inference is compromised. We focus on linear SDFs mainly because the useless factor problem is well defined for this class of models. In addition, we choose to present our results for the distance metric introduced by Hansen and Jagannathan (HJ, 1997). This measure has gained increased popularity in the empirical asset-pricing literature and has been used both as a model diagnostic and as a tool for model selection by many researchers.

The impact of the violation of this identification condition on the asymptotic properties of parameter restriction and specification tests in linear asset-pricing models estimated via generalized method of moments (GMM) was first studied by Kan and Zhang (1999a).<sup>1</sup> Kan and Zhang (1999a) analyze the behavior of the standard Wald test (which uses a variance matrix derived under the assumption of correct model specification) for potentially misspecified models. They show

---

<sup>1</sup> Burnside (2010, 2011) discusses analogous identification failures for alternative normalizations of the SDF. Kan and Zhang (1999b) study the consequences of lack of identification for two-pass cross-sectional regressions, while Kleibergen (2009, 2010) and Khalaf and Schaller (2011) propose test procedures that exhibit robustness to the degree of correlation between returns and factors in a two-pass cross-sectional regression framework.

that in this setup, using the standard Wald tests would lead one to conclude too often that the useless factor is priced. Furthermore, the specification tests have low power in rejecting a misspecified model. An immediate implication of this result is that many poor models and risk factors may have erroneously been deemed empirically successful, as our empirical applications illustrate.

We extend the analysis of Kan and Zhang (1999a) along several dimensions. First, unlike Kan and Zhang (1999a), we study the asymptotic and finite-sample properties of misspecification-robust parameter tests and investigate whether the model misspecification adjustment can restore the validity of the standard inference in the presence of useless factors. In particular, we demonstrate that the misspecification-robust Wald test for the significance of the SDF parameter on the useless factor is asymptotically distributed as a chi-squared random variable with one degree of freedom. This result is new to the literature and is somewhat surprising given the identification failure. It stands in sharp contrast with the Wald test constructed under the assumption of correct specification, which is shown by Kan and Zhang (1999a) to be asymptotically chi-squared distributed with degrees of freedom given by the difference between the number of assets and the number of factors included in the model. As a consequence, using standard inference will result in a rather extreme overrejection (with limiting rejection probability equal to one) of the null hypothesis that the risk premium on the useless factor is equal to zero.<sup>2</sup>

Second, we add to the analysis in Kan and Zhang (1999a) by also studying the limiting behavior of the estimates and Wald tests associated with the useful factors. We show that in misspecified models, the estimator of the coefficient associated with the useless factor diverges with the sample size, while the parameters on the useful factors are not consistently estimable. The limiting distributions of the  $t$ -statistics corresponding to the useful factors are found to be non-standard and less dispersed when a useless factor is present. Regardless of whether the model is correctly specified or misspecified, the misspecification-robust standard errors ensure asymptotically valid inference and allow us to identify factors that do not contribute to the pricing of the test assets. To conserve space, we relegate some of the theoretical results on the explicit form of the limiting distributions of the estimators, the  $t$ -tests under correct model specification and misspecification, as well as the HJ-distance test to an online appendix available on the authors' websites.

Third, we provide a constructive solution to the useless factor problem that restores the standard inference for the  $t$ -tests on the parameters associated with the useful factors and for the test of correct model specification. In particular, we propose an easy-to-implement sequential procedure that allows us to eliminate the useless factors from the model and show its asymptotic validity. More

---

<sup>2</sup> Our use of the term "overrejection" is somewhat nonstandard since the true risk premium on a useless factor is not identifiable. Nevertheless, since a useless factor does not improve the pricing performance of the model, testing the null of a zero risk premium is of most practical importance.

specifically, our model selection procedure is designed to remove factors that do not improve the model's pricing ability—that is, useless factors as well as useful factors with zero SDF parameters. In this paper, we refer to the useless and unpriced useful factors that do not reduce the HJ-distance as irrelevant factors. Monte Carlo simulation results suggest that our sequential model selection procedure is effective in retaining useful factors in the model and eliminating irrelevant factors. As a result, our proposed method is robust to both model misspecification and presence of useless factors in the analysis.<sup>3</sup>

Several remarks regarding our theoretical results are in order. We should stress that, similarly to White (1982) in a maximum likelihood framework, our misspecification-robust approach to inference allows for the model to be correctly specified and is asymptotically valid (albeit possibly slightly conservative) even when the model holds. This is important because a pre-test for correct model specification lacks power in distinguishing between correctly specified and misspecified models when a useless factor is included in the model. This leaves the misspecification-robust approach as the only feasible way to conduct inference, especially if a reduced rank test suggests an identification failure of the model. Another important issue that requires some clarification concerns our definition of a useless factor. While the paper studies the knife-edge case of a factor that is independent of the returns on the test assets, in practice all factors exhibit some nonzero correlation in finite samples and are probably better characterized as near-useless. Fortunately, our theoretical results conveniently encompass the possibility of weakly correlated factors without the need of local asymptotics, which is often used in the instrumental variable and unit root literature. It should be stressed that the presence of a useless factor represents the least favorable scenario for our proposed testing and model selection procedure based on the  $N(0, 1)$  critical values. In other words, our robust method is most conservative in the knife-edge case of a useless factor in the sense that it tends to underreject the null for the remaining useful factors. If the useless factor is instead replaced by a factor that exhibits a nonzero correlation with the returns, these underrejections will be smaller and the finite-sample properties of our method improve as we move away from the benchmark useless factor parameterization. We further discuss this issue in the simulation section of the paper.

Empirically, our interest is in robust estimation of several prominent asset-pricing models with macroeconomic and financial factors, also studied in Kan, Robotti, and Shanken (2013), using the HJ-distance measure. In addition to the basic CAPM and consumption CAPM (CCAPM), the theory-based models considered in our main empirical analysis are the CCAPM conditioned on the

---

<sup>3</sup> While we study explicitly only the GMM estimator based on the HJ-distance, our results continue to hold for the class of optimal GMM estimators. Some simulation results for the optimal GMM case are provided in Section 3 of the online appendix.

consumption-wealth ratio (CC-CAY) of Lettau and Ludvigson (2001), a time-varying version of the CAPM with human capital (C-LAB) of Jagannathan and Wang (1996), where the state variable driving the time variation in the SDF coefficients is the consumption-wealth ratio, the durable consumption model (D-CCAPM) of Yogo (2006), and the five-factor implementation of the intertemporal CAPM (ICAPM) used by Petkova (2006). We also study the well-known “three-factor model” of Fama and French (FF3, 1993). Although this model was primarily motivated by empirical observation, its size and book-to-market factors are sometimes viewed as proxies for more fundamental economic factors.

Our main empirical analysis uses the one-month T-bill, the monthly gross returns on the 25 Fama-French size and book-to-market portfolios, and the monthly gross returns on the 17 Fama-French industry portfolios from February 1959 until December 2012. The industry portfolios are included to provide a greater challenge to the various asset-pricing models, as recommended by Lewellen, Nagel, and Shanken (2010). The HJ-distance test rejects the hypothesis of correct specification for all models. In addition, the test for reduced rank indicates that only CAPM and FF3, two models with traded factors only, are properly identified. This clearly points to the need for statistical methods that are robust to model misspecification and weak identification. We show empirically that when misspecification-robust standard errors are employed, several macroeconomic factors – notably, the durable and nondurable consumption factors, the consumption-wealth factor of Lettau and Ludvigson (2001) and its interaction with nondurable consumption, labor income and the market return, the term and default spreads in ICAPM – do not appear to be priced at the 5% significance level. The only factors that survive our sequential procedure, which eliminates useless factors and the factors with zero risk premia, are the market factor in CAPM and FF3 and the book-to-market factor in FF3.

It is important to stress that the useless factor problem is not an isolated problem limited to the data and asset-pricing models considered in our main empirical analysis. We show that qualitatively similar pricing conclusions can be reached using different test assets and SDF specifications. Overall, our results suggest that the statistical evidence on the pricing ability of many macroeconomic and financial factors is weak, and their usefulness in explaining the cross-section of asset returns should be interpreted with caution.

The rest of the paper is organized as follows. Section 1 reviews some of the main results for asymptotically valid inference under potential model misspecification. In Section 2, we introduce a useless factor in the analysis and present limiting results for the parameters of interest and their  $t$ -statistics under both correct model specification and model misspecification. In Section 3, we discuss some practical implications of our theoretical analysis and suggest an easy-to-implement and asymptotically valid model selection procedure. Section 4 reports results from a Monte Carlo simulation experiment. In

Section 5, we investigate the performance of some popular asset-pricing models with traded and nontraded factors. Section 6 concludes.

### 1. Asymptotic Inference with Useful Factors

This section introduces the notation and reviews some main results that will be used in the subsequent analysis. Let

$$y_t(\gamma_1) = \tilde{f}_t' \gamma_1 \tag{1}$$

be a candidate linear SDF, where  $\tilde{f}_t = [1, f_t']$  is a  $K$ -vector with  $f_t$  being a  $(K - 1)$ -vector of risk factors, and  $\gamma_1$  is a  $K$ -vector of SDF parameters with generic element  $\gamma_{1i}$  for  $i = 1, \dots, K$ . The specification in Equation (1) is general enough to allow  $\tilde{f}_t$  to include cross-product terms (using lagged state variables as scaling factors); see Cochrane (1996).

Also, let  $x_t$  be the random payoffs of  $N$  assets at time  $t$  and  $q \neq 0_N$  be a vector of their original costs. This setup covers the case of gross returns on the test assets. For the case of excess returns ( $q = 0_N$ ), the mean of the SDF cannot be identified and researchers have to choose some normalization of the SDF (see, for example, Kan and Robotti 2008 and Burnside 2010). The theoretical and simulation results for the case of excess returns are very similar to those of the gross returns case presented below and are provided in Section 2 of the online appendix. We assume throughout that the second moment matrix of  $x_t$ ,  $U = E[x_t x_t']$ , is nonsingular so that none of the test assets are redundant.

Define the model pricing errors as

$$e(\gamma_1) = E[x_t \tilde{f}_t' \gamma_1 - q] = B \gamma_1 - q, \tag{2}$$

where  $B = E[x_t \tilde{f}_t']$ . If there exists no value of  $\gamma_1$  for which  $e(\gamma_1) = 0_N$ , the model is misspecified. This corresponds to the case when  $q$  is not in the span of the column space of  $B$ . The pseudo-true parameter vector  $\gamma_1^*$  is defined as the solution to the quadratic minimization problem

$$\gamma_1^* = \underset{\gamma_1 \in \Gamma_1}{\operatorname{argmin}} e(\gamma_1)' W e(\gamma_1) \tag{3}$$

for some symmetric and positive-definite weighting matrix  $W$ , where  $\Gamma_1$  denotes the parameter space (see also White 1982 for a definition of a pseudo-true value in a maximum likelihood framework).

The HJ-distance is obtained when  $W = U^{-1}$  and is given by

$$\delta = \sqrt{e(\gamma_1^*)' U^{-1} e(\gamma_1^*)}. \tag{4}$$

Given the computational simplicity and the nice economic and maximum pricing error interpretation of the HJ-distance, this measure of model misspecification is often used in applied work for estimation and evaluation of asset-pricing models. For this reason, we consider explicitly only the case

of the HJ-distance, although results for the optimal GMM estimator are also available from the authors upon request.

The estimator  $\tilde{\gamma}_1$  of  $\gamma_1^*$  is obtained by minimizing the sample analog of Equation (3):

$$\tilde{\gamma}_1 = \arg \min_{\gamma_1 \in \Gamma_1} \hat{e}(\gamma_1)' \hat{U}^{-1} \hat{e}(\gamma_1), \tag{5}$$

where  $\hat{U} = \frac{1}{T} \sum_{t=1}^T x_t x_t'$ ,  $\hat{e}(\gamma_1) = \hat{B} \gamma_1 - q$  and

$$\hat{B} = \frac{1}{T} \sum_{t=1}^T x_t f_t'. \tag{6}$$

Then, the solution to the above minimization problem is given by

$$\tilde{\gamma}_1 = (\hat{B}' \hat{U}^{-1} \hat{B})^{-1} \hat{B}' \hat{U}^{-1} q. \tag{7}$$

Let  $e_t(\gamma_1^*) = x_t f_t' \gamma_1^* - q$  and  $S = E[e_t(\gamma_1^*) e_t(\gamma_1^*)']$ . Assuming that  $[x_t', f_t']'$  are jointly stationary and ergodic processes with finite fourth moments,  $e_t(\gamma_1^*) - e(\gamma_1^*)$  forms a martingale difference sequence and  $B$  is of full column rank, Kan and Robotti (2009) show that

$$\sqrt{T}(\tilde{\gamma}_1 - \gamma_1^*) \xrightarrow{d} N(0_K, \Sigma_{\tilde{\gamma}_1}), \tag{8}$$

where  $\Sigma_{\tilde{\gamma}_1} = E[h_t h_t']$ ,

$$h_t = (B' U^{-1} B)^{-1} B' U^{-1} e_t(\gamma_1^*) + (B' U^{-1} B)^{-1} (f_t' - B' U^{-1} x_t) u_t \tag{9}$$

and

$$u_t = e(\gamma_1^*)' U^{-1} x_t. \tag{10}$$

Note that if the model is correctly specified (i.e.,  $u_t = 0$ ), the expression for  $h_t$  specializes to

$$h_t^0 = (B' U^{-1} B)^{-1} B' U^{-1} e_t(\gamma_1^*) \tag{11}$$

and the asymptotic covariance matrix of  $\sqrt{T}(\tilde{\gamma}_1 - \gamma_1^*)$  is simplified to

$$\Sigma_{\tilde{\gamma}_1}^0 = E[h_t^0 h_t^{0'}] = (B' U^{-1} B)^{-1} B' U^{-1} S U^{-1} B (B' U^{-1} B)^{-1}. \tag{12}$$

Suppose now that the interest lies in testing hypotheses on the individual parameters of the form  $H_0: \gamma_{1i} = \gamma_{1i}^*$  (for  $i = 1, \dots, K$ ) and define a selector vector  $\mathbf{t}_i$  with one for its  $i$ -th element and zero otherwise (the length of  $\mathbf{t}_i$  is implied by the matrix that it is multiplied to). Then, the  $t$ -statistic for  $\tilde{\gamma}_{1i}$  with standard error computed under potential model misspecification is asymptotically distributed as

$$t_m(\tilde{\gamma}_{1i}) = \frac{\tilde{\gamma}_{1i} - \gamma_{1i}^*}{\sqrt{\mathbf{t}_i' \hat{\Sigma}_{\tilde{\gamma}_1} \mathbf{t}_i}} \xrightarrow{d} N(0, 1), \tag{13}$$

where  $\hat{\Sigma}_{\tilde{\gamma}_1}$  is a consistent estimator of  $\Sigma_{\tilde{\gamma}_1}$ . Note that this result is valid irrespective of whether the model is misspecified or correctly specified.

In applied work, it is a common practice to test parameter restrictions using  $t$ -tests based on standard errors computed under the assumption of correct model specification. For this reason, it is instructive to consider the large sample behavior of the  $t$ -test

$$t_c(\tilde{\gamma}_{1i}) = \frac{\tilde{\gamma}_{1i} - \gamma_{1i}^*}{\sqrt{\mathbf{t}'_i \hat{\Sigma}_{\tilde{\gamma}_1}^0 \mathbf{t}_i}}, \tag{14}$$

where  $\hat{\Sigma}_{\tilde{\gamma}_1}^0$  is a consistent estimator of  $\Sigma_{\tilde{\gamma}_1}^0$ . If the model is indeed correctly specified, the  $t$ -test  $t_c(\tilde{\gamma}_{1i})$  is asymptotically distributed as a standard normal random variable

$$t_c(\tilde{\gamma}_{1i}) \xrightarrow{d} N(0, 1). \tag{15}$$

However, using the result in Equations (8)–(9), we have that under misspecified models

$$t_c(\tilde{\gamma}_{1i}) \xrightarrow{d} N\left(0, \frac{\mathbf{t}'_i \Sigma_{\tilde{\gamma}_1} \mathbf{t}_i}{\mathbf{t}'_i \Sigma_{\tilde{\gamma}_1}^0 \mathbf{t}_i}\right). \tag{16}$$

Furthermore, under the assumption that  $x_t$  and  $f_t$  are multivariate elliptically distributed, it can be shown (Kan and Robotti 2009) that  $(\mathbf{t}'_i \Sigma_{\tilde{\gamma}_1} \mathbf{t}_i) / (\mathbf{t}'_i \Sigma_{\tilde{\gamma}_1}^0 \mathbf{t}_i) > 1$ , which implies that standard inference based on critical values from the  $N(0, 1)$  distribution would tend to overreject the null hypothesis  $H_0: \gamma_{1i} = \gamma_{1i}^*$ . This also suggests that the testing procedure based on  $t_c$  would reject too often the null hypothesis that the SDF parameter is equal to zero.

We conclude this section with several observations that emerge from a closer inspection of the function  $h_t$  in Equation (9), which is used for computing the covariance matrix  $\Sigma_{\tilde{\gamma}_1}$  under misspecification. It proves useful to rewrite  $h_t$  as

$$h_t = h_t^0 + (B'U^{-1}B)^{-1}(\tilde{f}_t - B'U^{-1}x_t)u_t. \tag{17}$$

The adjustment term  $(B'U^{-1}B)^{-1}(\tilde{f}_t - B'U^{-1}x_t)u_t$  contains three components: (i) a misspecification component  $u_t$ , (ii) a spanning component  $\tilde{f}_t - B'U^{-1}x_t$  that measures the degree to which the factors are mimicked by the returns on the test assets, and (iii) a component  $(B'U^{-1}B)^{-1}$  that measures the usefulness of factors. The adjustment term is zero if the model is correctly specified ( $u_t = 0$ ) or if the factors are fully mimicked by the returns ( $\tilde{f}_t = B'U^{-1}x_t$ ). If the factors are nearly uncorrelated with the returns (i.e.,  $B$  is close to zero), the component  $(B'U^{-1}B)^{-1}$  can be large and the adjustment term tends to dominate the behavior of  $h_t$ . Intuitively, near-uncorrelatedness between the factors and the returns magnifies the consequences of small model specification errors and imperfectly mimicked factors.

## 2. Asymptotic Inference in the Presence of a Useless Factor

As argued in the introduction, many popular asset-pricing models include macroeconomic risk factors that often exhibit very low correlations with the

returns on the test assets. For this reason, we now consider a candidate SDF that is given by

$$y_t = \tilde{f}'_t \gamma_1 + g_t \gamma_2, \tag{18}$$

where  $g_t$  is assumed to be a useless factor such that it is independent of  $x_t$  and  $\tilde{f}_t$  for all time periods. Note that the independence between  $g_t$  and  $x_t$  implies  $d = E[x_t g_t] = 0_N$ .

Now let  $D = [B, d]$ ,  $\gamma = [\gamma'_1, \gamma'_2]'$ ,  $e(\gamma) = D\gamma - q$ ,  $\hat{d} = \frac{1}{T} \sum_{t=1}^T x_t g_t$ , and  $\hat{D} = [\hat{B}, \hat{d}]$ . Note that since  $d = 0_N$ , the vector of pricing errors

$$e(\gamma) = B\gamma_1 + d\gamma_2 - q = B\gamma_1 - q \tag{19}$$

is independent of the choice of  $\gamma_2$ . For the pseudo-true values of the SDF parameters, we can set  $\gamma_1^*$  as in Equation (3), but the parameter associated with the useless factor ( $\gamma_2^*$ ) cannot be identified. In the following, we set  $\gamma_2^* = 0$ , which is a natural choice because we show that  $\hat{\gamma}_2$  is symmetrically distributed around zero (see Section 1 of the online appendix for details). While the pseudo-true value of  $\gamma_2^*$  is not identifiable, the sample estimates of the SDF parameters are always identified and are given by

$$\hat{\gamma} = (\hat{D}' \hat{U}^{-1} \hat{D})^{-1} \hat{D}' \hat{U}^{-1} q. \tag{20}$$

We make the following assumptions.

**Assumption 1.** Assume that (i)  $N > K + 1$ ; (ii)  $[x'_t, \tilde{f}'_t, g_t]'$  are jointly stationary and ergodic processes with finite fourth moments; (iii)  $e_t(\gamma_1^*) - e(\gamma_1^*)$  forms a martingale difference sequence; and (iv) the matrices  $B$  ( $N \times K$ ) and  $D$  ( $N \times (K + 1)$ ) have a column rank  $K$ .

**Assumption 2.** Let  $\epsilon_t = x_t - B(E[\tilde{f}_t \tilde{f}'_t])^{-1} \tilde{f}_t$  and assume that  $E[\epsilon_t \epsilon'_t | \tilde{f}_t] = \Sigma$  (conditional homoscedasticity).

Assumption 1 imposes relatively mild restrictions on the data. The martingale difference sequence assumption allows for time-varying second- and higher-order moments of the pricing errors. This martingale difference sequence structure can be further relaxed to allow for serially correlated  $e_t(\gamma_1^*) - e(\gamma_1^*)$  at the cost of more tedious notation. Part (iv) of Assumption 1 accommodates our setup in which  $g_t$  is a useless factor. Assumption 2 is made for convenience in order to simplify some limiting results and can also be relaxed.

Our first results are concerned with the limiting behavior of  $\hat{\gamma}_1$  and  $\hat{\gamma}_2$  under correctly specified and misspecified models. Before we present these results, note that  $D$  is not of full column rank due to the presence of a useless factor. Therefore, the identification of the parameter vector  $\gamma^*$  fails and the sufficient conditions for the consistency and the asymptotic normality of the GMM estimator are not satisfied.

**Proposition 1.** Assume that the conditions in Assumption 1 are satisfied.

- (a) If  $\delta = 0$ , that is, the model is correctly specified, we have  $\hat{\gamma}_1 - \gamma_1^* = O_p(T^{-1/2})$  and  $\hat{\gamma}_2 = O_p(1)$ .
- (b) If  $\delta > 0$ , that is, the model is misspecified, we have  $\hat{\gamma}_1 - \gamma_1^* = O_p(1)$  and  $\hat{\gamma}_2 = O_p(T^{1/2})$ .

The explicit forms of the limiting distributions of the estimators and the proof of Proposition 1 are available in Section 1 of the online appendix. All of the asymptotic distributions are non-normal, and only the rate of convergence for  $\hat{\gamma}_1$  under correctly specified models is standard. The estimator  $\hat{\gamma}_2$  for the parameter on the useless factor converges to a bounded random variable and, hence, it is inconsistent.<sup>4</sup> The presence of model misspecification further exacerbates the inference problems, with the estimator  $\hat{\gamma}_1$  becoming inconsistent and the estimator  $\hat{\gamma}_2$  diverging at rate  $T^{1/2}$ .

Despite the highly non-standard limits of the SDF parameter estimates, it is possible that their  $t$ -statistics are well behaved. To investigate this, we define two types of  $t$ -statistics: (i)  $t_c(\hat{\gamma}_{1i})$ , for  $i = 1, \dots, K$ , and  $t_c(\hat{\gamma}_2)$  that use standard errors obtained under the assumption that the model is correctly specified, and (ii)  $t_m(\hat{\gamma}_{1i})$ , for  $i = 1, \dots, K$ , and  $t_m(\hat{\gamma}_2)$  that use standard errors under potentially misspecified models. The two types of  $t$ -statistics are based on the estimated covariance matrices  $\hat{\Sigma}_{\hat{\gamma}}^0 = \frac{1}{T} \sum_{t=1}^T \hat{h}_t^0 \hat{h}_t^{0'}$  and  $\hat{\Sigma}_{\hat{\gamma}} = \frac{1}{T} \sum_{t=1}^T \hat{h}_t \hat{h}_t'$ , where

$$\hat{h}_t^0 = (\hat{D}' \hat{U}^{-1} \hat{D})^{-1} \hat{D}' \hat{U}^{-1} \hat{e}_t, \tag{21}$$

$$\hat{h}_t = \hat{h}_t^0 + (\hat{D}' \hat{U}^{-1} \hat{D})^{-1} ([\tilde{f}_t', g_t']' - \hat{D}' \hat{U}^{-1} x_t) \hat{e}' \hat{U}^{-1} x_t, \tag{22}$$

$\hat{e}_t = x_t' (\tilde{f}_t' \hat{\gamma}_1 + g_t \hat{\gamma}_2) - q$  and  $\hat{e} = \frac{1}{T} \sum_{t=1}^T \hat{e}_t$ . We explicitly consider the behavior of  $t_c(\hat{\gamma}_{1i})$  and  $t_c(\hat{\gamma}_2)$  because it is a common practice for researchers to assume correct specification when computing the  $t$ -statistics.

In particular, the  $t$ -statistics of  $H_0: \gamma_{1i} = \gamma_{1i}^*$  and  $H_0: \gamma_2 = 0$  under the assumption of a correctly specified model have the form

$$t_c(\hat{\gamma}_{1i}) = \frac{\sqrt{T}(\hat{\gamma}_{1i} - \gamma_{1i}^*)}{\sqrt{t_i' \hat{\Sigma}_{\hat{\gamma}}^0 t_i}} \tag{23}$$

and

$$t_c(\hat{\gamma}_2) = \frac{\sqrt{T} \hat{\gamma}_2}{\sqrt{t_{K+1}' \hat{\Sigma}_{\hat{\gamma}}^0 t_{K+1}}}. \tag{24}$$

Kan and Zhang (1999a) studied the limiting behavior of  $t_c(\hat{\gamma}_2)$ , when no useful factor is present in the model, and showed that under  $H_0: \gamma_2 = 0$ ,  $t_c(\hat{\gamma}_2)^2$  is

<sup>4</sup> Convergence to a random variable arises in other contexts, such as spurious regressions with nonstationary data and instrumental variable models with weak instruments.

stochastically dominated by  $\chi_1^2$  for correctly specified models and  $t_c(\hat{\gamma}_2)^2 \xrightarrow{d} \chi_{N-K}^2$  for misspecified models. As stated below, these results continue to hold in the presence of useful factors. We further extend the results in Kan and Zhang (1999a) by deriving the explicit form of the limiting distributions of  $t_c(\hat{\gamma}_2)$  for correctly specified models and of  $t_c(\hat{\gamma}_{1i})$  for correctly specified and misspecified models (see Section 1 of the online appendix), which allows us to compute the limiting rejection probabilities of these tests when  $N(0, 1)$  critical values are used for inference.

The  $t$ -statistics of  $H_0 : \gamma_{1i} = \gamma_{1i}^*$  and  $H_0 : \gamma_2 = 0$  under a potentially misspecified model are given by

$$t_m(\hat{\gamma}_{1i}) = \frac{\sqrt{T}(\hat{\gamma}_{1i} - \gamma_{1i}^*)}{\sqrt{\iota_i' \hat{\Sigma}_{\hat{\gamma}} \iota_i}} \tag{25}$$

and

$$t_m(\hat{\gamma}_2) = \frac{\sqrt{T} \hat{\gamma}_2}{\sqrt{\iota_{K+1}' \hat{\Sigma}_{\hat{\gamma}} \iota_{K+1}}} \tag{26}$$

One of the main contributions of this paper is to establish that the use of misspecification robust standard errors in constructing the  $t$ -tests restores the standard inference on  $\gamma_2$  in misspecified models. In the other three cases (inference on  $\gamma_2$  in correctly specified models or on  $\gamma_1$  in correctly specified and misspecified models), the misspecification-robust  $t$ -tests are still asymptotically valid, but they tend to be conservative. The results for  $t_m(\hat{\gamma}_{1i})$  and  $t_m(\hat{\gamma}_2)$  are presented in the following proposition. For completeness, we also include the results for  $t_c(\hat{\gamma}_{1i})$  and  $t_c(\hat{\gamma}_2)$  under correctly specified and misspecified models. For presentational convenience, Proposition 2 states the limiting results for the squared  $t$ -tests (Wald tests).

**Proposition 2.**

- (a) Suppose that the conditions in Assumptions 1 and 2 hold. If  $\delta = 0$ , that is, the model is correctly specified, then  $t_c(\hat{\gamma}_{1i})^2$ ,  $t_c(\hat{\gamma}_2)^2$ ,  $t_m(\hat{\gamma}_{1i})^2$ , and  $t_m(\hat{\gamma}_2)^2$  are stochastically dominated by  $\chi_1^2$ .
- (b) Suppose that the conditions in Assumption 1 hold. If  $\delta > 0$ , that is, the model is misspecified, then  $t_c(\hat{\gamma}_{1i})^2$  and  $t_m(\hat{\gamma}_{1i})^2$  are stochastically dominated by  $\chi_1^2$ , and


$$t_c(\hat{\gamma}_2)^2 \xrightarrow{d} \chi_{N-K}^2, \tag{27}$$

$$t_m(\hat{\gamma}_2)^2 \xrightarrow{d} \chi_1^2. \tag{28}$$

Explicit expressions for the limiting distributions of the four tests statistics in part (a) and  $t_c(\hat{\gamma}_{1i})$  and  $t_c(\hat{\gamma}_2)$  in part (b) as well as the proof of Proposition 2 are provided in Section 1 of the online appendix. Proposition 2 illustrates the implications of using standard inference procedures (critical values from

$N(0, 1)$  for testing the individual statistical significance of the SDF parameters  $\gamma$  in the presence of a useless factor. Apart from  $t_m(\hat{\gamma}_2)$  in misspecified models, all the other statistics are not asymptotically distributed as standard normal random variables. For example, in misspecified models, the test statistic  $t_c(\hat{\gamma}_2)$  will overreject the null hypothesis when  $N(0, 1)$  is used as a reference distribution, and this overrejection increases with the number of test assets  $N$  (see Equation (27)). As a result, researchers will conclude erroneously (with high probability) that the factor  $g_t$  is important and should be included in the model. In order to visualize the source of the overrejection problem, Figure 1 plots the probability density function of  $t_c(\hat{\gamma}_2)$  for  $N - K = 7$  when the model is misspecified (see Theorem 2 in the online appendix).

Given the bimodal shape and a variance of 7 for the limiting distribution of  $t_c(\hat{\gamma}_2)$ , using the critical values from the standard normal distribution would obviously result in highly misleading inference. Importantly, part (b) of Proposition 2 shows that the  $t$ -statistic under potentially misspecified models,  $t_m(\hat{\gamma}_2)$ , retains its standard normal asymptotic distribution even when the factor is useless, and Figure 1 provides a graphical illustration of this result. The reduction in the degrees of freedom from  $N - K$  for the asymptotic chi-squared



**Figure 1**  
**Asymptotic distributions of  $t_c(\hat{\gamma}_2)$  and  $t_m(\hat{\gamma}_2)$  under misspecified models**

The figure presents the probability density functions of the limiting distributions of  $t_c(\hat{\gamma}_2)$  and  $t_m(\hat{\gamma}_2)$ , the  $t$ -statistics for the useless factor that use standard errors constructed under correctly specified and potentially misspecified models, respectively, for  $N - K = 7$  (see part (b) of Theorem 2 in the online appendix).

distribution of  $t_c(\hat{\gamma}_2)^2$  to 1 for the asymptotic chi-squared distribution of  $t_m(\hat{\gamma}_2)^2$  is striking.

Proposition 2 also suggests that the presence of a useless factor renders the inference on all the remaining parameters non-standard. Testing the statistical significance of the parameters on the useful factors, in both correctly specified and misspecified models, against the standard normal critical values would lead to underrejection of the null hypothesis and conservative inference.

The main conclusion that emerges from these results is that one should use misspecification-robust  $t$ -statistics when testing the statistical significance of individual SDF parameters. This will ensure that the statistical decision from this test is robust to possible model misspecification and useless factors. If the model happens to be correctly specified, this will result in conservative inference, but the useless factor will be removed with probability greater than  $1 - \alpha$ , where  $\alpha$  is the size of the test. If a useless factor is not present in the model, the standard normal asymptotics for the misspecification-robust test is restored as discussed in Section 1.<sup>5</sup>

### 3. Model Selection Procedure

It is worthwhile stressing an important aspect of the approach adopted in this paper. Economic theory often dictates which risk factors should be included in the model. In a regression framework, the inclusion of economic factors that are irrelevant does not affect the statistical inference in the model and only results in slightly inflated standard errors. In contrast, in the SDF setup studied in this paper, the inclusion of a useless factor has detrimental effects on the statistical inference on the remaining SDF parameters, their associated  $t$ -statistics and the model specification test. Our theoretical analysis reveals that the presence of a useless factor renders the remaining parameter estimates inconsistent and causes their  $t$ -statistics under both correct model specification and model misspecification to underreject the null. Only after the useless factor is identified and removed using the misspecification-robust  $t$ -test are the validity of the inference and the consistency of the parameters restored.

These considerations suggest that a sequential procedure based on the misspecification-robust  $t$ -statistics is necessary. One possibility is to select a subset of risk factors that survive a sequential testing procedure of individual significance of the SDF parameters  $\gamma$  based on  $p$ -values that are obtained from the quantiles of the  $N(0, 1)$  distribution and compared with a common significance level  $\alpha$ . Specifically, in the first stage, the full model is estimated

<sup>5</sup> Theorem 3 in the online appendix characterizes the limiting behavior of the sample squared HJ-distance in the presence of a useless factor and reveals that the HJ-distance test of correct model specification is inconsistent under the alternative.

and the factor with the smallest (in absolute value)  $t$ -statistic, for which the null of zero risk premium is not rejected at the prespecified nominal level, is eliminated from the model. The model is then reestimated with only the factors that survive the first stage. This procedure is repeated until either all factors are eliminated or the SDF parameter estimates on the remaining factors are found to be statistically significant at the desired nominal level when using the misspecification-robust  $t$ -test.

However, this model selection procedure does not account for the multitude of tests and could result in a substantially inflated rate of false discoveries (i.e., falsely identifying risk factors as being priced) depending on the number of tested hypotheses, the number of irrelevant factors and the dependence structure of the individual tests. The common solution to this multiple testing problem is to devise adjustments in the testing procedure that control the probability of one or more false discoveries, which is referred to in the statistical literature as familywise error rate. For a review of the large literature on controlling the familywise error rate in multiple testing problems and model selection, see Romano, Shaikh, and Wolf (2008), and for some recent applications of these techniques in empirical finance, see Barras, Scaillet, and Wermers (2010) and Bajgrowicz and Scaillet (2012). In this paper, we adopt the Bonferroni method for controlling the number of false discoveries and ensuring the asymptotic validity of our model selection procedure. While several refinements of the Bonferroni procedure have been proposed in the literature (see Benjamini and Hochberg 1995; Goeman and Solari 2010; Romano, Shaikh, and Wolf 2008; among others), the Bonferroni method has some advantages that are appealing to applied researchers.

First, the Bonferroni modification to the multiple testing problem is rather straightforward. Let  $K$  be the total number of SDF parameters being tested and let the  $p$ -value for testing the hypothesis  $H_0: \gamma_i = 0$  be denoted by  $p_{T,i}$ . Recall that all of the hypotheses  $H_0: \gamma_i = 0$  for  $i = 1, \dots, K$  are tested by comparing the misspecification-robust  $t$ -statistic against the critical value of the  $N(0, 1)$  distribution for a significance level  $\alpha$ . Then, the Bonferroni method rejects the null hypothesis if  $p_{T,i} \leq \alpha/K$  instead of  $p_{T,i} \leq \alpha$  as in the standard approach. Second, while some of the proposed methods require independence or some knowledge of the dependence structure of the individual test statistics, the Bonferroni method is robust to any form of dependence by assuming a worst-case dependence structure (Romano, Shaikh, and Wolf 2008). Naturally, the cost of this robustness is conservative inference. However, given that the number of risk factors in asset-pricing models rarely exceeds five, we show in our simulations that the Bonferroni method is only mildly conservative (i.e., it tends to select the relevant factors with high probability) while it fully controls the familywise error rate.

The Bonferroni method is asymptotically valid if (Romano, Shaikh, and Wolf 2008)

$$\limsup_{T \rightarrow \infty} P\{p_{T,i} \leq u\} \leq u,$$

for any uniformly distributed random variable  $u$  on the interval  $(0, 1)$ . The asymptotic validity of our proposed model selection procedure then follows from combining this asymptotic control of the Bonferroni method with the result in Proposition 2 that states that the limiting distribution of  $t_m(\hat{\gamma}_i)^2$  is stochastically dominated by the  $\chi_1^2$  distribution.

Instead of eliminating factors with insignificant  $t$ -ratios one at a time, one may be tempted to drop all the factors with insignificant  $t$ -ratios in each stage. Unlike our proposed method, this alternative procedure can lead to the undesirable outcome of eliminating multiple useful factors when a linear combination of them is useless.<sup>6</sup> In this situation, only one of these useful factors should be dropped to restore the full rank condition for the remaining factors. The effectiveness of our model selection procedure in eliminating useless factors (and factors with zero risk premia) and retaining useful factors in the model is analyzed in the simulation section below.

#### 4. Monte Carlo Simulations

In this section, we undertake a Monte Carlo experiment to assess the small-sample properties of the various test statistics in models with useful and useless factors. In our simulations, we also evaluate the effectiveness of the sequential model selection procedure described above in retaining useful factors and eliminating useless factors and factors with zero risk premia.

##### 4.1 Size of tests of parameter restrictions

For the analysis of the SDF parameter restriction tests, we consider three linear models: (i) a model with a constant term and a useful factor, (ii) a model with a constant term and a useless factor, and (iii) a model with a constant term, a useful factor, and a useless factor. For each model, we consider two separate cases: the case in which the model is correctly specified and the case in which the model is misspecified. The returns on the test assets, denoted by  $R_t$ , and the useful factor  $f_t$  are drawn from a multivariate normal distribution with mean  $\mu$  and covariance matrix  $V$ , where

$$\mu = E \begin{bmatrix} f_t \\ R_t \end{bmatrix} = \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix} \tag{29}$$

and

$$V = \text{Var} \begin{bmatrix} f_t \\ R_t \end{bmatrix} = \begin{bmatrix} V_{11} & V_{12} \\ V_{21} & V_{22} \end{bmatrix}. \tag{30}$$

Let  $\hat{\mu}$  and  $\hat{V}$  denote the sample estimates obtained from actual data. In all simulation designs, the covariance matrix of the simulated test asset returns  $V_{22}$

<sup>6</sup> See Section 1 of the online appendix for a simulation experiment in which no single factor is useless but a linear combination of them is.

is set equal to the estimated covariance matrix  $\hat{V}_{22}$  from the 1959:2–2012:12 sample of monthly gross returns on the one-month T-bill, the 25 Fama-French size and book-to-market ranked portfolios, and the 17 Fama-French industry portfolios (from Kenneth French’s website). Note that in this case,  $N = 43$  and  $q = 1_N$ . For the simulated useful factor, we calibrate its mean  $\mu_1$  and variance  $V_{11}$  to the sample mean  $\hat{\mu}_1$  and sample variance  $\hat{V}_{11}$  of the value-weighted market excess return. The covariances between the useful factor and the returns,  $V_{12}$ , are chosen based on the covariances estimated from the data,  $\hat{V}_{12}$ .

The pseudo-true values of the SDF parameters on the constant and the useful factor,  $\gamma^* = [\gamma_0^*, \gamma_1^*]'$ , are chosen as

$$\gamma^* = (X' \hat{V}_{22}^{-1} X)^{-1} X' \hat{V}_{22}^{-1} 1_N,$$

where  $X = [\hat{\mu}_2, \hat{V}_{21} + \hat{\mu}_2 \hat{\mu}_1]$ . Since the pricing errors of the model are given by

$$e(\gamma^*) = E[R_t(\gamma_0^* + f_t \gamma_1^*)] - 1_N = \mu_2(\gamma_0^* + \mu_1 \gamma_1^*) + V_{21} \gamma_1^* - 1_N, \quad (31)$$

we set  $\mu_2 = (1_N - V_{21} \gamma_1^*) / (\gamma_0^* + \mu_1 \gamma_1^*)$  for the pricing errors to be zero and the model to be correctly specified. For misspecified models, the means of the simulated returns,  $\mu_2$ , are set equal to the sample means of the actual returns  $\hat{\mu}_2$ . Finally, the useless factor is generated as a standard normal random variable independent of the returns and the useful factor. Using this approach to generating data allows the models in our simulation experiment to exhibit a similar degree of misspecification as some benchmark asset-pricing models such as the CAPM. More specifically, the HJ-distance for models (i) and (iii) is 0.523, while the HJ-distance for model (ii) is 0.535.

The time-series sample sizes that we consider in the simulations are  $T = 200$ , 600, and 1000. These choices of  $T$  cover the range of sample sizes that are typically encountered in empirical work with quarterly ( $T = 200$ ) and monthly ( $T = 600$ ) data. The sample size  $T = 1000$  is used to assess the quality of our asymptotic approximations. We also present the limiting rejection probabilities based on our asymptotic results in Theorem 2 in the online appendix.<sup>7</sup>

In Tables 1 to 3, we report the probabilities of rejection (based on 100,000 simulations) of  $H_0: \gamma_i = \gamma_i^*$  for models (i), (ii), and (iii), respectively, where the  $\gamma_i^*$ ’s for the constant and the useful factor are the chosen pseudo SDF parameters, and the  $\gamma_i^*$  for the useless factor is set equal to zero. We present results by comparing two different  $t$ -statistics with the standard normal distribution, the one computed under the assumption that the model is correctly specified,  $t_c(\hat{\gamma}_i)$ , and the one computed under the assumption that the model is potentially misspecified,  $t_m(\hat{\gamma}_i)$ . For each table, Panel A reports the probabilities of rejection when the model is correctly specified, and Panel B reports the probabilities of rejection when the model is misspecified.

<sup>7</sup> The limiting rejection probabilities of  $t_c$  in Table 1, Panel B, are computed based on Equation (16) assuming that the factor and the returns are multivariate normally distributed.

**Table 1**  
**Empirical size of the  $t$ -tests in a model with a useful factor**

$t$ -test	$T$	$\hat{\gamma}_0$			$\hat{\gamma}_1$		
		10%	5%	1%	10%	5%	1%
Panel A: Correctly specified model							
$t_c$	200	0.168	0.139	0.108	0.098	0.049	0.010
	600	0.134	0.098	0.063	0.099	0.049	0.009
	1,000	0.123	0.081	0.044	0.101	0.049	0.009
	$\infty$	0.100	0.050	0.010	0.100	0.050	0.010
$t_m$	200	0.168	0.139	0.108	0.097	0.049	0.010
	600	0.134	0.098	0.063	0.099	0.049	0.009
	1,000	0.123	0.081	0.044	0.101	0.049	0.009
	$\infty$	0.100	0.050	0.010	0.100	0.050	0.010
Panel B: Misspecified model							
$t_c$	200	0.173	0.144	0.112	0.099	0.048	0.009
	600	0.137	0.099	0.063	0.098	0.049	0.010
	1000	0.123	0.081	0.043	0.098	0.048	0.010
	$\infty$	0.100	0.050	0.010	0.100	0.050	0.010
$t_m$	200	0.173	0.144	0.112	0.099	0.048	0.009
	600	0.137	0.099	0.063	0.098	0.049	0.010
	1000	0.123	0.081	0.043	0.098	0.048	0.010
	$\infty$	0.100	0.050	0.010	0.100	0.050	0.010

The table presents the empirical size of the  $t$ -tests of  $H_0: \gamma_i = \gamma_i^*$  ( $i=0, 1$ ) in a model with a constant and a useful factor.  $\gamma_0$  is the coefficient on the constant term, and  $\gamma_1$  is the coefficient on the useful factor.  $t_c$  denotes the  $t$ -test constructed under the assumption of correct model specification, and  $t_m$  denotes the misspecification-robust  $t$ -test. We report results for different levels of significance (10%, 5%, and 1%) and for different values of the number of time-series observations ( $T$ ) using 100,000 simulations, assuming that the returns are generated from a multivariate normal distribution with means and covariance matrix calibrated to the 25 size and book-to-market Fama-French portfolio returns, the 17 Fama-French industry portfolio returns, and the one-month T-bill rate for the period 1959:2–2012:12. The various  $t$ -statistics are compared with the critical values from a standard normal distribution.

The results in Table 1, Panel A, show that for models that are correctly specified and contain only useful factors, the standard asymptotics provides an accurate approximation of the finite-sample behavior of the  $t$ -tests. Since the useful factor, calibrated to the properties of the value-weighted market excess return, is closely replicated by the returns on the test assets, the differences between the  $t$ -tests under correctly specified models ( $t_c$ ) and the  $t$ -tests under potentially misspecified models ( $t_m$ ) exhibit no differences even when the model fails to hold exactly (see Panel B).

Tables 2 and 3 present the empirical size of the  $t$ -tests in the presence of a useless factor. The simulation results for the  $t$ -tests on the parameters of the useful factor (and the constant term) confirm our theoretical findings that the null hypothesis is underrejected when  $N(0, 1)$  is used as a reference distribution. This is the case for correctly specified and misspecified models. For example, the underrejections are particularly pronounced for the  $t_m$  test on the constant term in Tables 2 and 3.

Similarly, the inference on the useless factor proves to be conservative when the model is correctly specified. However, when the model is misspecified, there are substantial differences between  $t_c$  and  $t_m$  for the useless factor. Under this scenario, we argued in Section 2 that the  $t$ -statistics under correct

**Table 2**  
**Empirical size of the  $t$ -tests in a model with a useless factor**

$t$ -test	$T$	$\hat{\gamma}_0$			$\hat{\gamma}_1$		
		10%	5%	1%	10%	5%	1%
Panel A: Correctly specified model							
$t_c$	200	0.008	0.002	0.000	0.131	0.067	0.012
	600	0.002	0.000	0.000	0.099	0.046	0.006
	1000	0.002	0.000	0.000	0.098	0.045	0.007
	$\infty$	0.001	0.000	0.000	0.088	0.039	0.005
$t_m$	200	0.001	0.000	0.000	0.036	0.012	0.001
	600	0.000	0.000	0.000	0.022	0.006	0.000
	1000	0.000	0.000	0.000	0.022	0.006	0.000
	$\infty$	0.000	0.000	0.000	0.018	0.004	0.000
Panel B: Misspecified model							
$t_c$	200	0.020	0.006	0.000	0.328	0.234	0.103
	600	0.020	0.005	0.000	0.472	0.383	0.228
	1000	0.024	0.007	0.000	0.557	0.476	0.327
	$\infty$	0.088	0.039	0.005	1.000	1.000	1.000
$t_m$	200	0.002	0.000	0.000	0.081	0.036	0.005
	600	0.001	0.000	0.000	0.081	0.038	0.006
	1000	0.001	0.000	0.000	0.086	0.041	0.007
	$\infty$	0.001	0.000	0.000	0.100	0.050	0.010

The table presents the empirical size of the  $t$ -tests of  $H_0 : \gamma_i = \gamma_i^*$  ( $i = 0, 1$ ) in a model with a constant and a useless factor.  $\gamma_0$  is the coefficient on the constant term, and  $\gamma_1$  is the coefficient on the useless factor.  $t_c$  denotes the  $t$ -test constructed under the assumption of correct model specification, and  $t_m$  denotes the misspecification-robust  $t$ -test. We report results for different levels of significance (10%, 5%, and 1%) and for different values of the number of time-series observations ( $T$ ) using 100,000 simulations, assuming that the returns are generated from a multivariate normal distribution with means and covariance matrix calibrated to the 25 size and book-to-market Fama-French portfolio returns, the 17 Fama-French industry portfolio returns, and the one-month T-bill rate for the period 1959:2–2012:12. The various  $t$ -statistics are compared with the critical values from a standard normal distribution. The rejection rates for the limiting case ( $T = \infty$ ) are based on the asymptotic distributions given in Theorem 2 in the online appendix.

model specification have a non-normal asymptotic distribution, while the misspecification-robust  $t$ -statistic for the parameter on the useless factor has a  $N(0, 1)$  asymptotic distribution. Since the  $t_c$  test on the useless factor is asymptotically distributed (up to a sign) as  $\sqrt{\chi_{N-K}^2}$ , it tends to overreject severely when the critical values from  $N(0, 1)$  are used, and the degree of overrejection increases with the sample size. In contrast, the  $t_m$  test on the useless factor has good size properties, although for small sample sizes, it slightly underrejects. As the sample size increases, the empirical rejection rates approach the limiting rejection probabilities (as shown in the rows for  $T = \infty$ ) computed from the corresponding asymptotic distributions in Theorem 2 in the online appendix.

Some comments about the underrejections reported in Tables 2 and 3 for useful (in correctly specified and misspecified models) and useless (for correctly specified models) factors are in order. Starting with the useless factor, it turns out that the statistical underrejection of the null hypothesis of a zero SDF coefficient is actually economically beneficial for our selection procedure since this useless factor will be selected even less often than the nominal size of the test. For the useful factor, the underrejections may raise more concerns. However, note that

**Table 3**  
**Empirical size of the  $t$ -tests in a model with a useful and a useless factor**

$t$ -test	$T$	$\hat{\gamma}_0$			$\hat{\gamma}_1$			$\hat{\gamma}_2$		
		10%	5%	1%	10%	5%	1%	10%	5%	1%
Panel A: Correctly specified model										
$t_c$	200	0.054	0.023	0.007	0.094	0.045	0.008	0.128	0.066	0.012
	600	0.059	0.028	0.008	0.096	0.047	0.009	0.100	0.046	0.006
	1000	0.056	0.025	0.007	0.096	0.046	0.009	0.095	0.043	0.005
	$\infty$	0.052	0.020	0.002	0.096	0.047	0.009	0.088	0.039	0.005
$t_m$	200	0.027	0.011	0.004	0.090	0.042	0.007	0.036	0.012	0.001
	600	0.037	0.016	0.006	0.092	0.045	0.008	0.022	0.006	0.000
	1000	0.037	0.016	0.005	0.092	0.044	0.008	0.019	0.005	0.000
	$\infty$	0.037	0.014	0.002	0.092	0.045	0.008	0.018	0.004	0.000
Panel B: Misspecified model										
$t_c$	200	0.056	0.023	0.005	0.095	0.046	0.008	0.319	0.227	0.098
	600	0.057	0.023	0.004	0.095	0.046	0.009	0.464	0.376	0.224
	1000	0.057	0.021	0.003	0.094	0.046	0.009	0.550	0.469	0.319
	$\infty$	0.088	0.039	0.005	0.088	0.039	0.005	1.000	1.000	1.000
$t_m$	200	0.016	0.006	0.002	0.086	0.040	0.006	0.080	0.036	0.005
	600	0.013	0.005	0.002	0.079	0.037	0.006	0.082	0.038	0.006
	1000	0.009	0.003	0.001	0.071	0.032	0.005	0.087	0.042	0.007
	$\infty$	0.001	0.000	0.000	0.001	0.000	0.000	0.100	0.050	0.010

The table presents the empirical size of the  $t$ -tests of  $H_0: \gamma_i = \gamma_i^*$  ( $i=0, 1, 2$ ) in a model with a constant, a useful, and a useless factor.  $\gamma_0$  is the coefficient on the constant term,  $\gamma_1$  is the coefficient on the useful factor, and  $\gamma_2$  is the coefficient on the useless factor.  $t_c$  denotes the  $t$ -test constructed under the assumption of correct model specification, and  $t_m$  denotes the misspecification-robust  $t$ -test. We report results for different levels of significance (10%, 5%, and 1%) and for different values of the number of time-series observations ( $T$ ) using 100,000 simulations, assuming that the returns are generated from a multivariate normal distribution with means and covariance matrix calibrated to the 25 size and book-to-market Fama-French normal portfolio returns, the 17 Fama-French industry portfolio returns, and the one-month T-bill rate for the period 1959:2–2012:12. The various  $t$ -tests are compared with the critical values from a standard normal distribution. The rejection rates for the limiting case ( $T = \infty$ ) are based on the asymptotic distributions given in Theorem 2 in the online appendix.

the  $t$ -tests for the constant and the useful factors in Tables 1 to 3 are testing the null hypothesis  $H_0: \gamma_i = \gamma_i^*$ . In practice, the researchers are interested in testing the null hypothesis of whether the factor is priced or not—that is,  $H_0: \gamma_i = 0$ . While the underrejections in Tables 2 and 3 will affect the local power of the test, it is still quite possible that these tests reject with reasonably high probability the null  $H_0: \gamma_i = 0$  for useful factors with a nonzero risk premium parameter. This is illustrated in the model selection procedure below, which selects factors based on whether their parameter estimates are significantly different from zero or not. The results show that the useful factors with nonzero risk premium parameters tend to be selected with sufficiently high probability (97–99% for  $T = 1000$ ).

As mentioned in the introduction, all factors are expected to exhibit a nonzero correlation with the returns in finite samples, and the case of a completely useless factor may seem extreme. We reiterate, however, that the useless factor case represents the least favorable scenario in our testing setup, as it places an upper bound on the conservativeness of our robust method. In Section 1 of the online appendix, we numerically illustrate this by considering a model with a priced useful factor and an unpriced factor, where the unpriced factor is

calibrated to three observed factors with a different correlation (weak, moderate, and strong) with the returns on the test assets. The simulation results clearly indicate that the misspecification-robust  $t$ -test for the priced useful factor exhibits smaller underrejections (and improved power) compared with those in Table 3, while the rejection rates for the unpriced factor remain largely unchanged. The complete set of results and the specifics of the simulation design are presented in Section 1 of the online appendix.

#### 4.2 Survival rates of risk factors in the sequential testing procedure

Our findings suggest that the misspecification-robust  $t$ -test should be used when it is uncertain whether the factor is useful or useless and the model is correctly specified or misspecified. However, since the procedure based on the  $t_m$  test is often conservative, some useful factors might be erroneously excluded from the model. The frequency at which this happens is evaluated in the model selection procedure presented below.

Table 4 reports the survival rates of different factors when using the sequential procedure described in Section 3. In particular, we compare the survival rates from using the  $t_m$  test with the survival rates from using the  $t_c$  test. The false discovery rate of the multiple testing problem is controlled using the Bonferroni method. In our simulations, we consider (i) a linear model with a constant term, two useful factors with  $\gamma_i^* \neq 0$ , an unpriced useful factor with  $\gamma_i^* = 0$ , and a useless factor, and (ii) a linear model with a constant term, two useful factors with  $\gamma_i^* \neq 0$ , and two useless factors. The mean and variance of the useful factors with  $\gamma_i^* \neq 0$  are calibrated to the mean and variance of the excess market return and the high-minus-low factor of Fama and French (1993). The mean and variance of the unpriced useful factor with  $\gamma_i^* = 0$  are calibrated to the mean and variance of the small-minus-big factor of Fama and French (1993). As in Tables 1–3, the returns and the useful factors are drawn from a multivariate normal distribution.<sup>8</sup> Finally, the useless factor is generated as a standard normal random variable, independent of the test asset returns and the useful factors. The time-series sample size is taken to be  $T = 200, 600,$  and  $1000$ . The nominal level of the sequential testing procedure is set equal to 5%. The probability that at least one irrelevant (useless or unpriced) factor survives the model selection ( $MS$ ) procedure is reported in the last two columns of the table, with  $MS_c$  denoting the survival probability computed from the  $t_c$  tests and  $MS_m$  denoting the survival probability computed from the  $t_m$  tests.

Panel A shows that when the model is correctly specified, the procedures based on  $t_c$  and  $t_m$  do a similarly good job in retaining the useful factors with nonzero SDF parameters in the model (the survival probabilities are 99–100% for  $T = 600$ ) and eliminating the irrelevant factors. This indicates

<sup>8</sup> See Gospodinov, Kan, and Robotti (2013) for a description on how to impose zero restrictions on the parameters in correctly specified and misspecified models. The HJ-distance of the misspecified model used in this section is 0.522.

**Table 4**  
**Survival rates of risk factors: Two useful and two irrelevant factors**

Panel A: Correctly specified model

T	Useful ( $\gamma_1^* \neq 0$ )		Useful ( $\gamma_2^* \neq 0$ )		Useful ( $\gamma_3^* = 0$ )		Useless		Prob.	
	$t_c(\hat{\gamma}_1)$	$t_m(\hat{\gamma}_1)$	$t_c(\hat{\gamma}_2)$	$t_m(\hat{\gamma}_2)$	$t_c(\hat{\gamma}_3)$	$t_m(\hat{\gamma}_3)$	$t_c(\hat{\gamma}_4)$	$t_m(\hat{\gamma}_4)$	$MS_c$	$MS_m$
200	0.519	0.517	0.674	0.661	0.027	0.023	0.023	0.002	0.049	0.025
600	0.986	0.987	0.999	0.999	0.014	0.013	0.009	0.001	0.023	0.014
1000	1.000	1.000	1.000	1.000	0.013	0.013	0.008	0.000	0.022	0.013

T	Useful ( $\gamma_1^* \neq 0$ )		Useful ( $\gamma_2^* \neq 0$ )		Useless		Useless		Prob.	
	$t_c(\hat{\gamma}_1)$	$t_m(\hat{\gamma}_1)$	$t_c(\hat{\gamma}_2)$	$t_m(\hat{\gamma}_2)$	$t_c(\hat{\gamma}_3)$	$t_m(\hat{\gamma}_3)$	$t_c(\hat{\gamma}_4)$	$t_m(\hat{\gamma}_4)$	$MS_c$	$MS_m$
200	0.520	0.521	0.666	0.658	0.023	0.002	0.024	0.002	0.046	0.004
600	0.988	0.989	0.999	0.999	0.009	0.001	0.010	0.001	0.019	0.001
1000	1.000	1.000	1.000	1.000	0.008	0.000	0.008	0.000	0.016	0.001

Panel B: Misspecified model

T	Useful ( $\gamma_1^* \neq 0$ )		Useful ( $\gamma_2^* \neq 0$ )		Useful ( $\gamma_3^* = 0$ )		Useless		Prob.	
	$t_c(\hat{\gamma}_1)$	$t_m(\hat{\gamma}_1)$	$t_c(\hat{\gamma}_2)$	$t_m(\hat{\gamma}_2)$	$t_c(\hat{\gamma}_3)$	$t_m(\hat{\gamma}_3)$	$t_c(\hat{\gamma}_4)$	$t_m(\hat{\gamma}_4)$	$MS_c$	$MS_m$
200	0.498	0.502	0.648	0.633	0.033	0.025	0.108	0.008	0.138	0.033
600	0.971	0.981	0.993	0.996	0.018	0.014	0.206	0.008	0.220	0.022
1000	0.996	0.998	0.999	0.999	0.015	0.012	0.295	0.009	0.306	0.021

T	Useful ( $\gamma_1^* \neq 0$ )		Useful ( $\gamma_2^* \neq 0$ )		Useless		Useless		Prob.	
	$t_c(\hat{\gamma}_1)$	$t_m(\hat{\gamma}_1)$	$t_c(\hat{\gamma}_2)$	$t_m(\hat{\gamma}_2)$	$t_c(\hat{\gamma}_3)$	$t_m(\hat{\gamma}_3)$	$t_c(\hat{\gamma}_4)$	$t_m(\hat{\gamma}_4)$	$MS_c$	$MS_m$
200	0.300	0.308	0.398	0.389	0.105	0.007	0.106	0.007	0.205	0.015
600	0.853	0.900	0.921	0.951	0.202	0.009	0.204	0.009	0.384	0.018
1000	0.959	0.983	0.981	0.992	0.279	0.010	0.282	0.010	0.517	0.020

The table presents the survival rates of the factors in a model with a constant, two useful factors (with  $\gamma_1^* \neq 0$  and  $\gamma_2^* \neq 0$ ), and two irrelevant factors. The first irrelevant factor is either a useful factor that does not contribute to pricing (with  $\gamma_3^* = 0$ ) or a useless factor (with  $\gamma_3^*$  unidentified), and the second irrelevant factor is a useless factor (with  $\gamma_4^*$  unidentified). The sequential procedure is implemented by using the misspecification-robust  $t$ -tests ( $t_m(\hat{\gamma}_i)$  column) as well as the  $t$ -tests under correctly specified models ( $t_c(\hat{\gamma}_i)$  column). The false discovery rate of the multiple testing procedure is controlled using the Bonferroni method. The last two columns of the table report the probability that at least one useless or unpriced useful factor survives using the  $t$ -tests under correctly specified models ( $MS_c$ ) and misspecification-robust  $t$ -tests ( $MS_m$ ). The nominal level of the sequential testing procedure is set equal to 5%. We report results for different values of the number of time-series observations ( $T$ ) using 100,000 simulations, assuming that the returns are generated from a multivariate normal distribution with means and covariance matrix calibrated to the 25 size and book-to-market Fama-French portfolio returns, the 17 Fama-French industry portfolio returns, and the one-month T-bill rate for the period 1959:2–2012:12.

that using the  $t_c$  test in the presence of a useless factor is not problematic when the underlying model holds exactly. However, as shown in Panel B, the situation drastically changes when the model is misspecified. In this case, the procedures based on  $t_c$  and  $t_m$  still retain the useful factors with nonzero SDF parameters with similarly high probability (85–100% for  $T=600$ ), but they produce very different results when it comes to the useless factor. For example, despite its conservative nature (due to the Bonferroni adjustment), the procedure based on  $t_c$  will retain the useless factor 28–30% of the time for  $T=1000$ . In contrast, the procedure based on  $t_m$  will retain the useless factor only about 0.8–0.9% of the time for  $T=1000$ . Similarly, the probability of at least one irrelevant factor being selected in the final specification of the model is 31–52% (2.0–2.1%) for  $T=1000$  when the  $t_c$  ( $t_m$ ) test is used and the model

is misspecified. It should be emphasized that the effectiveness of the proposed sequential procedure in retaining the useful factors in the model depends on the correlation between the useful factors and the returns on the test assets and on the magnitude of the SDF coefficient associated with the useful factor. Our procedure will be more effective in retaining a useful factor in the model, the higher this correlation and the larger the SDF coefficient on the useful factor.

Some aspects of our model selection procedure deserve additional remarks. It should be stressed that the objective of our proposed sequential test is to find the most parsimonious model with the same HJ-distance as the full model. The fact that our method eliminates unpriced useful factors along with useless factors is not of particular concern since these factors do not contribute to the reduction of the HJ-distance. As a result, our model selection procedure purges the model from useless factors (that give rise to inference problems) and unpriced factors (that do not improve the pricing ability of the model) and restores the standard asymptotic theory for the remaining factors in the SDF.

## 5. Empirical Analysis

Our theoretical and simulation results point out some serious pitfalls in the empirical analysis of asset-pricing models with nontraded factors. In the following, we use monthly data to demonstrate the relevance of our theoretical results.

To show that our findings are not specific to the test assets and SDFs considered in the main empirical application, we also use an alternative set of test assets and SDFs with macroeconomic risk factors that are available only at quarterly frequency.

### 5.1 Main application

First, we describe the data used in the empirical analysis and outline the different specifications of the asset-pricing models considered. Then, we present our results.

**5.1.1 Data and asset-pricing models.** As in the Monte Carlo simulations, the test asset returns are the monthly gross returns on the one-month T-bill, the value-weighted 25 Fama-French size and book-to-market ranked portfolios, and the 17 industry portfolios from Kenneth French's website.<sup>9</sup> The data are from February 1959 until December 2012. We analyze seven asset-pricing models, starting with the simple static CAPM. The SDF specification for this model is

$$y_t^{CAPM}(\gamma) = \gamma_0 + \gamma_1 v w_t, \tag{32}$$

<sup>9</sup> Using only the 25 Fama-French size and book-to-market ranked portfolios as test assets yields qualitatively similar results.

where  $vw$  is the excess return (in excess of the one-month T-bill rate) on the value-weighted stock market index (NYSE-AMEX-NASDAQ) from Kenneth French's website. The CAPM performed well in the early tests—for example, Fama and MacBeth (1973)—but has fared poorly since.

One extension that has performed better is our second model, the three-factor model (FF3) of Fama and French (1993). This model extends the CAPM by including the two empirically motivated factors  $smb$  and  $hml$ , where  $smb$  is the return difference between portfolios of stocks with small and large market capitalizations, and  $hml$  is the return difference between portfolios of stocks with high and low book-to-market ratios (“value” and “growth” stocks, respectively) from Kenneth French's website. The SDF specification is

$$y_t^{FF3}(\gamma) = \gamma_0 + \gamma_1 vw_t + \gamma_2 smb_t + \gamma_3 hml_t. \quad (33)$$

The third model (ICAPM) is an empirical implementation of Merton's (1973) intertemporal extension of the CAPM based on Campbell (1996), who argues that innovations in state variables that forecast future investment opportunities should serve as factors. The five-factor specification proposed by Petkova (2006) is

$$y_t^{ICAPM}(\gamma) = \gamma_0 + \gamma_1 vw_t + \gamma_2 term_t + \gamma_3 def_t + \gamma_4 div_t + \gamma_5 rf_t, \quad (34)$$

where  $term$  is the difference between the yields of ten-year and one-year government bonds (from the Board of Governors of the Federal Reserve System),  $def$  is the difference between the yields of long-term corporate Baa bonds (from the Board of Governors of the Federal Reserve System) and long-term government bonds (from Ibbotson Associates),  $div$  is the dividend yield on the Center for Research in Security Prices (CRSP) value-weighted stock market portfolio, and  $rf$  is the one-month T-bill yield (from CRSP, Fama Risk Free Rates). The actual factors for  $term$ ,  $def$ ,  $div$ , and  $rf$  are their innovations from a VAR(4) system of seven state variables that also includes  $vw$ ,  $smb$ , and  $hml$ . Following Petkova (2006), the innovations are orthogonalized and scaled to have the same variance as the innovation in the excess market return.

Our fourth model is the scaled CAPM specification including human capital (C-LAB) of Jagannathan and Wang (1996), also considered by Lettau and Ludvigson (2001). This model incorporates measures of the return on human capital as well as the change in financial wealth, and allows the conditional SDF coefficients to vary with the consumption-aggregate wealth ratio ( $cay$ ) of Lettau and Ludvigson (2001).<sup>10</sup> The SDF specification is

$$y_t^{C-LAB}(\gamma) = \gamma_0 + \gamma_1 cay_{t-1} + \gamma_2 vw_t + \gamma_3 lab_t + \gamma_4 vw_t \cdot cay_{t-1} + \gamma_5 lab_t \cdot cay_{t-1}, \quad (35)$$

where  $lab$  is the growth rate in per capita labor income,  $L$ , defined as the difference between total personal income and dividend payments, divided

<sup>10</sup> Following Jørgensen–Vissing and Attanasio (2003), we linearly interpolate the quarterly values of  $cay$  to permit analysis at the monthly frequency.

by the total population (from the Bureau of Economic Analysis). Following Jagannathan and Wang (1996), we use a two-month moving average to construct the growth rate  $lab_t = (L_{t-1} + L_{t-2}) / (L_{t-2} + L_{t-3}) - 1$ , for the purpose of minimizing the influence of measurement error. The SDF specification in Equation (35) is obtained by scaling the constant term, the  $vw$  and the  $lab$  factors by a constant and  $cay$ . Scaling factors by instruments is one popular way of allowing factor risk premia to vary over time. Gagliardini, Ossola, and Scaillet (2013), for example, provide strong empirical evidence of time-varying risk premia in standard factor models.

Next, we consider consumption-based models. Our fifth model (CCAPM) is the unconditional consumption model with

$$y_t^{CCAPM}(\gamma) = \gamma_0 + \gamma_1 c_{nd,t}, \tag{36}$$

where  $c_{nd}$  is the growth rate in real per capita nondurable consumption (seasonally adjusted at annual rates) from the Bureau of Economic Analysis. This model has generally not performed well empirically (see Lettau and Ludvigson 2001 for a summary of the poor empirical performance of CCAPM). Therefore, we also examine two other consumption models that have yielded more encouraging results.

One such model (CC-CAY) is a conditional version of the CCAPM due to Lettau and Ludvigson (2001). The relation is

$$y_t^{CC-CAY}(\gamma) = \gamma_0 + \gamma_1 c_{nd,t} + \gamma_2 cay_{t-1} + \gamma_3 c_{nd,t} \cdot cay_{t-1}. \tag{37}$$

The SDF specification in Equation (37) is obtained by scaling the constant term and the  $c_{nd}$  factor by a constant and  $cay$ .

The last model (D-CCAPM), due to Yogo (2006), highlights the cyclical role of *durable* consumption in asset pricing. The specification is

$$y_t^{D-CCAPM}(\gamma) = \gamma_0 + \gamma_1 vw_t + \gamma_2 c_{nd,t} + \gamma_3 c_{d,t}, \tag{38}$$

where  $c_d$  is the growth rate in real per capita durable consumption (seasonally adjusted at annual rates) from the Bureau of Economic Analysis.

**5.1.2 Results.** Before presenting the estimation results for the SDF parameters, we first investigate whether the various risk factors are correlated with the asset returns and whether the seven models described above are properly identified. As mentioned in the theoretical section of the paper, the presence of a useless factor leads to a violation of the crucial identification condition that the  $N \times K$  matrix  $B = E[x_t \tilde{f}'_t]$  is of full column rank. Therefore, it is of interest to test if  $B$  is of (reduced) rank  $K - 1$ . Since  $\hat{B}$  is not invariant to rescaling of the data, we first perform a normalization on  $\hat{B}$ . Define  $\hat{\Theta} = \hat{U}^{-\frac{1}{2}} \hat{B} \hat{S}_f^{-\frac{1}{2}}$  and its corresponding population counterpart  $\Theta = U^{-\frac{1}{2}} B S_f^{-\frac{1}{2}}$ , where  $\hat{S}_f = \frac{1}{T} \sum_{t=1}^T \tilde{f}_t \tilde{f}'_t$  and  $S_f = E[\tilde{f}_t \tilde{f}'_t]$ . Note that  $\hat{\Theta}$  is invariant to rescaling of the data and  $\text{rank}(\Theta) = \text{rank}(B)$ .

Let  $\hat{\Pi}$  be a consistent estimator of the asymptotic covariance matrix of  $\sqrt{T}\text{vec}(\hat{\Theta} - \Theta)$ , where  $\text{vec}(\cdot)$  is the vec operator. Following Ratsimalahelo (2002) and Kleibergen and Paap (2006), we perform a singular value decomposition on  $\hat{\Theta}$  such that  $\hat{\Theta} = \tilde{U} \tilde{S} \tilde{V}'$ , where  $\tilde{U}' \tilde{U} = I_N$ ,  $\tilde{V}' \tilde{V} = I_K$ , and  $\tilde{S}$  is an  $N \times K$  matrix with the singular values of  $\hat{\Theta}$  in decreasing order on its diagonal. Let  $\tilde{U}_2$  be the last  $N - K + 1$  columns of  $\tilde{U}$ ,  $\tilde{V}_2$  be the last column of  $\tilde{V}$ , and

$$\tilde{\Pi} = (\tilde{V}_2' \otimes \tilde{U}_2') \hat{\Pi} (\tilde{V}_2 \otimes \tilde{U}_2). \tag{39}$$

Then, a test of  $H_0 : \text{rank}(\Theta) = \text{rank}(B) = K - 1$  is given by

$$\mathcal{W}^* = T \tilde{s}_K^2 \tilde{\Pi}^{11} \xrightarrow{d} \chi_{N-K+1}^2, \tag{40}$$

where  $\tilde{s}_K$  is the smallest singular value of  $\hat{\Theta}$ , and  $\tilde{\Pi}^{11}$  is the  $(1, 1)$  element of  $\tilde{\Pi}^{-1}$ .

Table 5, Panel A, reports the rank restriction test ( $\mathcal{W}^*$ ) and its  $p$ -value ( $p$ -val) of the null that  $E[x_{it}(1, f_{it})]$  has a column rank of one. The panel shows that we cannot reject the null of a column rank of one at the 5% significance level for seven out of 14 macroeconomic and financial factors. This finding suggests that several risk factors in ICAPM, C-LAB, CCAPM, CC-CAY, and D-CCAPM can be reasonably considered as useless and that our asymptotic results on useless factors are of practical importance. Panel B further shows that only CAPM and FF3 pass the test of full rank condition at the 5% nominal level. This is consistent with the fact that  $vw$ ,  $smb$ , and  $hml$  are highly correlated with the returns on the test assets, while most factors in the other models are not. Panel B also shows that no model passes the HJ-distance specification test at conventional levels of significance. Since the HJ-distance test has been shown to substantially overreject under the null in realistic simulation settings with many test assets, we also consider an alternative test of  $H_0 : \lambda = U^{-1}e = 0_N$  (which is equivalent to the test of  $H_0 : \delta = 0$ ). Gospodinov, Kan, and Robotti (2013) show that this alternative Lagrange multiplier ( $LM$ ) test has excellent size and power properties. The results in Panel B indicate that one would reach the same conclusions using the  $LM$  and HJ-distance tests. Therefore, the model rejections documented in Table 5, Panel B, do not seem to be driven by the finite-sample properties of the HJ-distance test. Overall, these empirical findings suggest that valid inference should account for the fact that some of the models are potentially misspecified and poorly identified.

Although the rank restriction test serves as a useful pre-test for possible identification problems, this test does not allow us to unambiguously identify which factor contributes to the identification failure of the model. In addition, this test does not address the question of which risk factors are important in explaining the cross-sectional differences in asset returns. Our misspecification-robust test of  $H_0 : \gamma_i = 0$  proves to be of critical importance in (i) providing the direction of the identification failure and (ii) allowing us to determine whether a given risk factor is priced. Panels C and D of Table 5 present the  $t$ -tests

**Table 5**  
**Monthly analysis of some popular linear asset-pricing models**

Panel A: Rank test for individual factors

Test	<i>vw</i>	<i>smb</i>	<i>hml</i>	<i>term</i>	<i>def</i>	<i>div</i>	<i>rf</i>	<i>cay</i>	<i>lab</i>	<i>vw·cay</i>	<i>lab·cay</i>	<i>c<sub>nd</sub></i>	<i>c<sub>nd</sub>·cay</i>	<i>c<sub>d</sub></i>
$\mathcal{W}^*$	205.7	199.9	189.7	47.1	89.2	61.1	44.8	54.0	54.8	83.1	49.7	54.6	43.0	58.6
<i>p</i> -val	0.000	0.000	0.000	0.270	0.000	0.029	0.354	0.101	0.088	0.000	0.193	0.092	0.430	0.046

Panel B: HJ-distance, Lagrange multiplier, and rank tests

Model	$\hat{\delta}$	<i>p</i> -val	<i>LM</i>	<i>p</i> -val	$\mathcal{W}^*$	<i>p</i> -val
CAPM	0.523	0.000	139.766	0.000	205.7	0.000
FF3	0.487	0.000	121.265	0.000	190.7	0.000
ICAPM	0.469	0.000	84.567	0.000	42.1	0.298
C-LAB	0.481	0.000	78.892	0.000	31.1	0.780
CCAPM	0.513	0.000	112.888	0.000	54.6	0.092
CC-CAY	0.484	0.000	81.523	0.000	40.9	0.430
D-CCAPM	0.510	0.000	114.966	0.000	46.4	0.225

Panel C: Model selection procedure using standard errors under correct model specification

Model	<i>vw</i>	<i>smb</i>	<i>hml</i>	<i>term</i>	<i>def</i>	<i>div</i>	<i>rf</i>	<i>cay</i>	<i>lab</i>	<i>vw·cay</i>	<i>lab·cay</i>	<i>c<sub>nd</sub></i>	<i>c<sub>nd</sub>·cay</i>	<i>c<sub>d</sub></i>
CAPM	<b>-2.64</b>													
FF3	-3.43	-0.57	-4.59											
	<b>-3.69</b>		<b>-4.62</b>											
ICAPM	-2.04			-2.70	0.50	1.86	0.85							
	-2.06			-2.93		1.80	0.72							
	-2.10			-3.76		1.88								
	-2.20			-3.77										
				<b>-3.74</b>										
C-LAB	-0.72							0.15	0.59	2.46	-2.22			
	-0.71								0.60	2.45	-3.36			
	-0.80									2.39	-3.43			
										2.38	-3.63			
											<b>-3.79</b>			
CCAPM												<b>-3.23</b>		
CC-CAY								-3.13				-3.05	0.97	
								<b>-3.24</b>				<b>-3.03</b>		
D-CCAPM	-1.12											-2.41		-0.67
	-1.14											-2.58		
												<b>-3.23</b>		

(continued)

**Table 5**  
**Continued**

Panel D: Model selection procedure using model misspecification-robust standard errors

Model	<i>vw</i>	<i>smb</i>	<i>hml</i>	<i>term</i>	<i>def</i>	<i>div</i>	<i>rf</i>	<i>cay</i>	<i>lab</i>	<i>vw·cay</i>	<i>lab·cay</i>	<i>c<sub>nd</sub></i>	<i>c<sub>nd</sub>·cay</i>	<i>c<sub>d</sub></i>
CAPM	<b>-2.64</b>													
FF3	-3.35	-0.53	-4.40											
	<b>-3.68</b>		<b>-4.45</b>											
ICAPM	-2.04			-1.86	0.36	1.36	0.62							
	-2.06			-2.13		1.38	0.52							
	-2.10			-2.61		1.42								
	-2.20			-2.60										
	-2.52													
C-LAB	-0.64							0.08	0.36	1.88	-1.09			
	-0.61								0.37	1.88	-1.81			
	-0.71									1.87	-1.80			
										1.89	-2.07			
											-1.98			
CCAPM												-1.75		
CC-CAY								-1.85				-1.88	0.47	
								-1.89				-1.84		
								-1.89						
D-CCAPM	-0.90											-1.22		-0.41
	-0.90											-1.27		
												-1.75		

The table presents the estimation and testing results of the seven asset-pricing models described in Section 5.1.1. The models are estimated using monthly gross returns on the 25 size and book-to-market Fama-French portfolios, the 17 Fama-French industry portfolios, and the one-month T-bill. The data are from 1959:2 until 2012:12. Panel A reports the rank restriction test ( $\mathcal{W}^*$ ) and its  $p$ -value ( $p$ -val) of the null that  $E[x_t(1, f_{it})]$  has a column rank of one. In Panel B, we report the sample HJ-distance ( $\hat{\delta}$ ), the Lagrange multiplier ( $LM$ ) test, and the rank restriction test ( $\mathcal{W}^*$ ) with the corresponding  $p$ -values ( $p$ -val) for each model. The  $t$ -tests of the model selection procedures based on the standard errors under correct model specification and model misspecification are in Panels C and D, respectively. We use boldface to highlight those cases in which the factors survive the model selection procedure at the 5% significance level using the Bonferroni adjustment.

under correct model specification and potential model misspecification as well as the model selection procedure described in Section 3. Using  $t$ -tests under correct specification, the results in Panel C suggest that the factors *term* in ICAPM, *lab·cay* in C-LAB, *cay* in CC-CAY, and  $c_{nd}$  in CCAPM, CC-CAY, and D-CCAPM survive the sequential testing procedure at the 5% significance level using the Bonferroni correction. However, given the violation of the full rank condition for these models, the standard normal distribution is not the appropriate reference distribution in this case. Since some of these factors are very weakly correlated with the returns on the test assets and effectively behave as useless factors, they tend to be included in the model much more often than they should (see the simulation results for the irrelevant factors in Panel B of Table 4). Therefore, the model selection procedure needs to be implemented using misspecification-robust  $t$ -tests.

Panel D shows that none of the nontraded factors listed above survives the sequential procedure based on misspecification-robust  $t$ -tests at the 5% significance level. Finally, for traded factors, we find strong evidence of pricing for the *vw* factor in CAPM and the *vw* and *hml* factors in FF3 in both Panels C and D.<sup>11</sup>

## 5.2 Additional empirical evidence

The results in Table 5 suggest that the statistical evidence on the pricing ability of several macroeconomic and financial factors is weak and their usefulness in explaining the cross-section of asset returns should be interpreted with caution. In this section, we further emphasize the importance of accounting for model misspecification and weak identification in empirical work. The following application uses an alternative set of test assets and SDFs that includes macroeconomic risk factors whose data are available only at quarterly frequency.

The test asset returns are the quarterly gross returns on the one-month T-bill, the value-weighted 6 Fama-French size and book-to-market ranked portfolios, the 17 industry portfolios, and the 10 momentum portfolios from Kenneth French's website. The sample period is from 1952:Q2 until 2012:Q4. We consider the following asset-pricing specifications: (i) the conditional CCAPM (CC-CAY) version of Lettau and Ludvigson (2001) with  $c_{nd}$ , *cay*, and  $c_{nd}cay$  as described in the previous section; (ii) the conditional CAPM (C-ML) of Santos and Veronesi (2006) with *vw* and *vw* scaled by the labor income-consumption ratio (*ml*) as risk factors; (iii) a version of the conditional consumption CAPM

<sup>11</sup> It is possible that the individual risk factors considered in this section do not capture adequately the risk incorporated in all of the macroeconomic data that is available to market participants. One approach to extract parsimoniously the common variation in macroeconomic variables is the factor analysis advocated by Stock and Watson (2002). In unreported results, we follow this approach and construct three orthogonal factors that summarize the dynamics of 130 U.S. macroeconomic time series for the period March 1960–December 2011. Using the same set of test assets, we find that no factor survives the misspecification-robust model selection procedure at the 5% significance level using the Bonferroni adjustment.

(CC-MY) proposed by Lustig and Van Nieuwerburgh (2005) with the housing collateral ratio ( $my$ ),  $c_{nd}$ , and the interaction term  $c_{nd}my$  as risk factors; and (iv) the sector investment model (SIM) of Li, Vassalou, and Xing (2006) with the log investment growth rates for households ( $i_h$ ), nonfinancial corporations ( $i_c$ ), and non-corporate sector ( $i_{nc}$ ) as risk factors. We include the CC-CAY model again since the original data for the  $cay$  factor are available at a quarterly frequency. These four models with nontraded factors have yielded encouraging results in cross-sectional asset pricing.<sup>12</sup>

The empirical results for quarterly data are reported in Table 6, with Panel A showing that for all factors except for  $vw$  and  $vw \cdot ml$ , we cannot reject the null that  $E[x_t(1, f_{it})]$  has a column rank of one at the 5% significance level. In addition, the results in Panel B indicate that we cannot reject the null of reduced rank for all models, except for C-ML, and that all models are rejected by the HJ-distance and  $LM$  specification tests. This clearly points to the need of statistical procedures that are robust to model misspecification and weak identification.

Panel D of Table 6 shows that all factors except for  $vw \cdot ml$  do not survive the model selection procedure based on the misspecification-robust  $t$ -test.<sup>13</sup> This stands in sharp contrast to the results in Panel C of Table 6, where the  $t$ -test under correctly specified models is employed. However, our theoretical and simulation analyses clearly showed that relying on the  $t$ -test under correct specification is grossly inappropriate when the underlying model is misspecified and the factors are very weakly correlated with the returns on the test assets. As one example, consider CC-MY. In the final stage of the model selection procedure in Panel C, both  $cg$  and  $my$  seem to be priced. On the contrary, no factor in CC-MY survives the model selection procedure based on misspecification-robust  $t$ -tests. Taken together, these results serve as a warning signal to researchers that are interested in estimating and analyzing SDF parameters on nontraded risk factors.

## 6. Conclusion

It is well known that asset returns are, at best, only weakly correlated with many macroeconomic factors. Nonetheless, researchers in finance have typically relied on inference methods that are not robust to weak identification and model misspecification when evaluating the incremental pricing ability of these factors. Our paper demonstrates that when a model is misspecified, the standard  $t$ -test of statistical significance will lead us to erroneously conclude, with high

<sup>12</sup> We also estimated the CAPM, FF3, CCAPM, and D-CCAPM models considered in the previous section using quarterly data, and the results are very similar to the ones for the monthly application.

<sup>13</sup> Although  $vw \cdot ml$  is in principle a nontraded factor, it is very highly correlated with the returns on the test assets. This is a situation in which, as discussed at the end of Section 1, the  $t$ -ratios under correctly specified and misspecified models are likely to deliver a similar answer.

**Table 6**  
**Quarterly analysis of some popular linear asset-pricing models**

Panel A: Rank test for individual factors

Test	<i>vw</i>	<i>vw·ml</i>	<i>c<sub>nd</sub></i>	<i>cay</i>	<i>c<sub>nd</sub>·cay</i>	<i>my</i>	<i>c<sub>nd</sub>·my</i>	<i>i<sub>h</sub></i>	<i>i<sub>c</sub></i>	<i>i<sub>nc</sub></i>
<i>W</i> *	92.2	92.5	46.0	43.9	33.6	39.0	39.6	42.4	33.5	28.1
<i>p</i> -val	0.000	0.000	0.066	0.098	0.438	0.218	0.199	0.128	0.444	0.708

Panel B: HJ-distance, Lagrange multiplier, and rank tests

Model	$\hat{\delta}$	<i>p</i> -val	<i>LM</i>	<i>p</i> -val	<i>W</i> *	<i>p</i> -val
C-ML	0.790	0.000	95.378	0.000	54.2	0.008
CC-CAY	0.761	0.000	73.471	0.000	29.8	0.527
CC-MY	0.751	0.002	86.353	0.000	40.8	0.111
SIM	0.776	0.000	87.339	0.000	24.3	0.800

Panel C: *t*-tests using standard errors under correct model specification

Model	<i>vw</i>	<i>vw·ml</i>	<i>c<sub>nd</sub></i>	<i>cay</i>	<i>c<sub>nd</sub>·cay</i>	<i>my</i>	<i>c<sub>nd</sub>·my</i>	<i>i<sub>h</sub></i>	<i>i<sub>c</sub></i>	<i>i<sub>nc</sub></i>
C-ML	1.91	-1.97								
		<b>-2.99</b>								
CC-CAY			-2.77	-2.17	0.51					
			-2.95	-2.33						
			<b>-3.36</b>							
CC-MY			-2.74			2.51	-0.99			
			<b>-2.91</b>			<b>2.41</b>				
SIM								-2.89	-0.59	0.26
								-3.04	-0.60	
								<b>-3.42</b>		

Panel D: *t*-tests using model misspecification-robust standard errors

Model	<i>vw</i>	<i>vw·ml</i>	<i>c<sub>nd</sub></i>	<i>cay</i>	<i>c<sub>nd</sub>·cay</i>	<i>my</i>	<i>c<sub>nd</sub>·my</i>	<i>i<sub>h</sub></i>	<i>i<sub>c</sub></i>	<i>i<sub>nc</sub></i>
C-ML	1.34	-1.38								
		<b>-2.97</b>								
CC-CAY			-1.80	-1.57	0.31					
			-1.92	-1.64						
			-1.99							
CC-MY			-1.88			1.64	-0.64			
			-1.97			1.51				
			-1.99							
SIM								-1.92	-0.38	0.15
								-2.04	-0.38	
								-2.27		

The table presents the estimation and testing results of the four asset-pricing models described in Section 5.2. The models are estimated using quarterly gross returns on the 6 size and book-to-market Fama-French portfolios, the 17 Fama-French industry portfolios, the 10 momentum portfolios, and the one-month T-bill. The data are from 1952:Q2 until 2012:Q4. Panel A reports the rank restriction test (*W*\*) and its *p*-value (*p*-val) of the null that  $E[x_t(1, f_{it})]$  has a column rank of one. In Panel B, we report the sample HJ-distance ( $\hat{\delta}$ ), the Lagrange multiplier (*LM*) test, and the rank restriction test (*W*\*) with the corresponding *p*-values (*p*-val) for each model. The *t*-tests of the model selection procedures based on the standard errors under correct model specification and model misspecification are in Panels C and D, respectively. We use boldface to highlight those cases in which the factors survive the model selection procedure at the 5% significance level using the Bonferroni adjustment.

probability, that a useless factor is relevant and should be included in the model. Importantly, we show that the *t*-test of statistical significance will be valid only if it is computed using misspecification-robust standard errors. Furthermore, we argue that the presence of a useless factor affects the inference on the remaining model parameters and the test of correct specification. In particular, when a useless factor is present in the model, the limiting distributions of the *t*-statistics

for the useful factors are nonstandard and the HJ-distance specification test is inconsistent.

In order to overcome these problems, we propose an easy-to-implement sequential model selection procedure based on misspecification-robust  $t$ -tests that restores the standard inference on the parameters of interest. We show via simulations that the proposed procedure is effective in eliminating useless factors as well as factors that do not improve the pricing ability of the model.

Finally, we employ our methodology to investigate the empirical performance of several prominent asset-pricing models with traded and nontraded factors. While the market factor and the book-to-market factor of Fama and French (1993) are often found to be priced, the statistical evidence on the pricing ability of many nontraded factors is rather weak when using the model selection procedure based on misspecification-robust  $t$ -tests.

## References

- Bajgrowicz, P., and O. Scaillet. 2012. Technical trading revisited: Persistence tests, transaction costs, and false discoveries. *Journal of Financial Economics* 106:473–91.
- Barras, L., O. Scaillet, and R. Wermers. 2010. False discoveries in mutual fund performance: Measuring luck in estimated alphas. *Journal of Finance* 65:179–216.
- Benjamini, Y., and Y. Hochberg. 1995. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B* 57:289–300.
- Burnside, C. A. 2010. Identification and inference in linear stochastic discount factor models. NBER working paper 16634.
- Burnside, C. A. 2011. The cross-section of foreign currency risk premia and consumption growth risk: Comment. *American Economic Review* 101:3456–76.
- Campbell, J. Y. 1996. Understanding risk and return. *Journal of Political Economy* 104:298–345.
- Cochrane, J. H. 1996. A cross-sectional test of an investment-based asset pricing model. *Journal of Political Economy* 104:572–621.
- Fama, E. F., and J. D. MacBeth. 1973. Risk, return, and equilibrium: Empirical tests. *Journal of Political Economy* 71:607–36.
- Fama, E. F., and K. R. French. 1993. Common risk factors in the returns on stocks and bonds. *Journal of Financial Economics* 33:3–56.
- Gagliardini, P., E. Ossola, and O. Scaillet. 2013. Time-varying risk premia in large cross-sectional equity datasets. Working Paper, University of Lugano.
- Goeman, J. J., and A. Solari. 2010. The sequential rejection principle of familywise error control. *Annals of Statistics* 38:3782–3810.
- Gospodinov, N., R. Kan, and C. Robotti. 2013. Chi-squared tests for evaluation and comparison of asset pricing models. *Journal of Econometrics* 173:108–25.
- Hansen, L. P., and R. Jagannathan. 1997. Assessing specification errors in stochastic discount factor models. *Journal of Finance* 52:557–90.
- Jagannathan, R., and Z. Wang. 1996. The conditional CAPM and the cross-section of expected returns. *Journal of Finance* 51:3–53.
- Jørgensen-Vissing, A., and O. P. Attanasio. 2003. Stock-market participation, intertemporal substitution, and risk-aversion. *American Economic Review Papers and Proceedings* 93:383–91.

- Kan, R., and C. Robotti. 2008. Specification tests of asset pricing models using excess returns. *Journal of Empirical Finance* 15:816–38.
- Kan, R., and C. Robotti. 2009. Model comparison using the Hansen-Jagannathan distance. *Review of Financial Studies* 22:3449–90.
- Kan, R., C. Robotti, and J. Shanken. 2013. Pricing model performance and the two-pass cross-sectional regression methodology. *Journal of Finance* 68:2617–49.
- Kan, R., and C. Zhang. 1999a. GMM tests of stochastic discount factor models with useless factors. *Journal of Financial Economics* 54:103–27.
- Kan, R., and C. Zhang. 1999b. Two-pass tests of asset pricing models with useless factors. *Journal of Finance* 54:204–35.
- Khalaf, L. A., and H. Schaller. 2011. How Fama-MacBeth can go wrong – and an informative solution. Working Paper, Carleton University.
- Kleibergen, F. 2009. Tests of risk premia in linear factor models. *Journal of Econometrics* 149:149–73.
- Kleibergen, F. 2010. Reality checks for and of factor pricing. Working Paper, Brown University.
- Kleibergen, F., and R. Paap. 2006. Generalized reduced rank tests using the singular value decomposition. *Journal of Econometrics* 133:97–126.
- Lettau, M., and S. C. Ludvigson. 2001. Resurrecting the (C)CAPM: A cross-sectional test when risk premia are time-varying. *Journal of Political Economy* 109:1238–87.
- Lewellen, J. W., S. Nagel, and J. Shanken. 2010. A skeptical appraisal of asset-pricing tests. *Journal of Financial Economics* 96:175–94.
- Li, Q., M. Vassalou, and Y. Xing. 2006. Sector investment growth rates and the cross-section of equity returns. *Journal of Business* 79:1637–65.
- Ludvigson, S. C. 2013. Advances in consumption-based asset pricing: Empirical tests. Chapter 12 in *Handbook of the economics of finance*, Vol. 2, 799–906. Ed. G. M. Constantinides, M. Harris, and R. M. Stulz. Amsterdam: North Holland.
- Lustig, H., and S. Van Nieuwerburgh. 2005. Housing collateral, consumption insurance, and risk premia: An empirical perspective. *Journal of Finance* 60:1167–1219.
- Merton, R. C. 1973. An intertemporal capital asset pricing model. *Econometrica* 41:867–87.
- Petkova, R. 2006. Do the Fama-French factors proxy for innovations in predictive variables? *Journal of Finance* 61:581–612.
- Ratsimalahelo, Z. 2002. Rank test based on matrix perturbation theory. Working Paper, University of Franche-Comté.
- Romano, J. P., A. M. Shaikh, and M. Wolf. 2008. Formalized data snooping based on generalized error rates. *Econometric Theory* 24:404–47.
- Santos, T., and P. Veronesi. 2006. Labor income and predictable stock returns. *Review of Financial Studies* 19:1–44.
- Stock, J. H., and M. W. Watson. 2002. Forecasting using principal components from a large number of predictors. *Journal of the American Statistical Association* 97:1167–79.
- White, H. 1982. Maximum likelihood estimation of misspecified models. *Econometrica* 50:1–25.
- Yogo, M. 2006. A consumption-based explanation of expected stock returns. *Journal of Finance* 61:539–80.